

# Discrete Random Variables: Joint PMFs, Conditioning and Independence

Berlin Chen

Department of Computer Science & Information Engineering  
National Taiwan Normal University

Reference:

- D. P. Bertsekas, J. N. Tsitsiklis, *Introduction to Probability*, Sections 2.5-2.7

# Motivation

- Given an experiment, e.g., a medical diagnosis
  - The results of blood test is modeled as numerical values of a random variable  $X$
  - The results of magnetic resonance imaging (MRI,核磁共振攝影) is also modeled as numerical values of a random variable  $Y$

We would like to consider probabilities of events involving simultaneously the numerical values of these two variables and to investigate their mutual couplings

$$\mathbf{P} \left( \{X = x\} \cap \{Y = y\} \right)?$$

# Joint PMF of Random Variables

- Let  $X$  and  $Y$  be random variables associated with the same experiment (also the same sample space and probability laws), the **joint PMF** of  $X$  and  $Y$  is defined by

$$p_{X,Y}(x,y) = \mathbf{P}(\{X=x\} \cap \{Y=y\}) = \mathbf{P}(X=x, Y=y)$$

- if event  $A$  is the set of all pairs  $(x,y)$  that have a certain property, then the probability of  $A$  can be calculated by

$$\mathbf{P}((X,Y) \in A) = \sum_{(x,y) \in A} p_{X,Y}(x,y)$$

- Namely,  $A$  can be specified in terms of  $X$  and  $Y$

## Marginal PMFs of Random Variables (1/2)

- The **PMFs** of random variables  $X$  and  $Y$  can be calculated from their **joint PMF**


$$p_X(x) = \sum_y p_{X,Y}(x,y), \quad p_Y(y) = \sum_x p_{X,Y}(x,y)$$

- $p_X(x)$  and  $p_Y(y)$  are often referred to as the **marginal PMFs**
- The above two equations can be verified by

$$\begin{aligned} p_X(x) &= \mathbf{P}(X=x) \\ &= \sum_y \mathbf{P}(X=x, Y=y) \\ &= \sum_y p_{X,Y}(x,y) \end{aligned}$$

## Marginal PMFs of Random Variables (2/2)

- Tabular Method:** Given the joint PMF of random variables  $X$  and  $Y$  is specified in a two-dimensional table, the marginal PMF of  $X$  or  $Y$  at a given value is obtained by adding the table entries along a corresponding column or row, respectively



## Functions of Multiple Random Variables (1/2)

- A function  $Z = g(X, Y)$  of the random variables  $X$  and  $Y$  defines another random variable. Its PMF can be calculated from the joint PMF  $p_{X,Y}$

$$p_Z(z) = \sum_{\{(x,y) | g(x,y)=z\}} p_{X,Y}(x,y)$$

- The expectation for a function of several random variables

$$\mathbf{E}[Z] = \mathbf{E}[g(X, Y)] = \sum_x \sum_y g(x, y) p_{X,Y}(x, y)$$

## Functions of Multiple Random Variables (2/2)

- If the function of several random variables is linear and of the form  $Z = g(X, Y) = aX + bY + c$

$$\mathbf{E}[Z] = a\mathbf{E}[X] + b\mathbf{E}[Y] + c$$

- How can we verify the above equation ?

# An Illustrative Example

- Given the random variables  $X$  and  $Y$  whose joint is given in the following figure, and a new random variable  $Z$  is defined by  $Z = X + 2Y$ , calculate  $\mathbf{E}[Z]$

– Method 1:

$$\mathbf{E}[X] = 1 \cdot \frac{3}{20} + 2 \cdot \frac{6}{20} + 3 \cdot \frac{8}{20} + 4 \cdot \frac{3}{20} = \frac{51}{20}$$

$$\mathbf{E}[Y] = 1 \cdot \frac{3}{20} + 2 \cdot \frac{7}{20} + 3 \cdot \frac{7}{20} + 4 \cdot \frac{3}{20} = \frac{50}{20}$$

$$\mathbf{E}[Z] = \mathbf{E}[X] + 2\mathbf{E}[Y] = \frac{51}{20} + 2 \cdot \frac{50}{20} = \frac{151}{20} = 7.55$$


– Method 2:

$$p_Z(z) = \sum_{\{(x,y) | x+2y=z\}} p_{X,Y}(x,y)$$

$$p_Z(3) = \frac{1}{20}, p_Z(4) = \frac{1}{20}, p_Z(5) = \frac{2}{20}, p_Z(6) = \frac{2}{20}$$

$$p_Z(7) = \frac{4}{20}, p_Z(8) = \frac{3}{20}, p_Z(9) = \frac{3}{20}, p_Z(10) = \frac{2}{20}$$

$$p_Z(11) = \frac{1}{20}, p_Z(12) = \frac{1}{20}$$



$$\begin{aligned} \therefore \mathbf{E}[Z] &= 3 \cdot \frac{1}{20} + 4 \cdot \frac{1}{20} + 5 \cdot \frac{2}{20} + 6 \cdot \frac{2}{20} \\ &\quad + 7 \cdot \frac{4}{20} + 8 \cdot \frac{3}{20} + 9 \cdot \frac{3}{20} + 10 \cdot \frac{2}{20} \\ &\quad + 11 \cdot \frac{1}{20} + 12 \cdot \frac{1}{20} = 7.55 \end{aligned}$$



## More than Two Random Variables (1/2)

- The joint PMF of three random variables  $X$ ,  $Y$  and  $Z$  is defined in analogy with the above as

$$p_{X,Y,Z}(x, y, z) = \mathbf{P}(X = x, Y = y, Z = z)$$

- The corresponding marginal PMFs

$$p_{X,Y}(x, y) = \sum_z p_{X,Y,Z}(x, y, z)$$

and

$$p_X(x) = \sum_y \sum_z p_{X,Y,Z}(x, y, z)$$

## More than Two Random Variables (2/2)

- The expectation for the function of random variables  $X$ ,  $Y$  and  $Z$

$$\mathbf{E}[g(X, Y, Z)] = \sum_x \sum_y \sum_z g(x, y, z) p_{X, Y, Z}(x, y, z)$$

- If the function is linear and has the form  $aX + bY + cZ + d$

$$\mathbf{E}[aX + bY + cZ + d] = aE[X] + bE[Y] + cE[Z] + d$$

- A generalization to more than three random variables

$$\mathbf{E}[a_1X_1 + a_2X_2 + \cdots + a_nX_n] = a_1E[X_1] + a_2E[X_2] + \cdots + a_nE[X_n]$$

## An Illustrative Example

- **Example 2.10. Mean of the Binomial.** Your probability class has 300 students and each student has probability  $1/3$  of getting an A, independently of any other student.
  - What is the mean of  $X$ , the number of students that get an A?

Let

$$X_i = \begin{cases} 1, & \text{if the } i\text{th student gets an A} \\ 0, & \text{otherwise} \end{cases}$$

$\Rightarrow X_1, X_2, \dots, X_{300}$  are bernoulli random variables with common mean  $p = 1/3$

Their sum  $X = X_1 + X_2 + \dots + X_{300}$  can be interpreted as a binomial random variable with parameters  $n$  ( $n = 300$ ) and  $p$  ( $p = 1/3$ ). That is,  $X$  is the number of success in  $n$  ( $n = 300$ ) independent trials

$$\therefore \mathbf{E}[X] = \mathbf{E}[X_1 + X_2 + \dots + X_{300}] = \sum_{i=1}^{300} \mathbf{E}[X_i] = 300 \cdot 1/3 = 100$$

# Conditioning

- Recall that conditional probability provides us with a way to reason about the outcome of an experiment, based on partial information
- In the same spirit, we can define **conditional PMFs**, given the occurrence of a certain event or given the value of another random variable

# Conditioning a Random Variable on an Event (1/2)

- The **conditional PMF** of a random variable  $X$ , conditioned on a particular event  $A$  with  $\mathbf{P}(A) > 0$ , is defined by (where  $X$  and  $A$  are associated with the same experiment)

$$P_{X|A}(x) = \mathbf{P}(X = x|A) = \frac{\mathbf{P}(\{X = x\} \cap A)}{\mathbf{P}(A)}$$

- Normalization Property


- Note that the events  $\mathbf{P}(\{X = x\} \cap A)$  are **disjoint** for different values of  $X$ , their union is  $A$

$$\mathbf{P}(A) = \sum_x \mathbf{P}(\{X = x\} \cap A) \quad \text{Total probability theorem}$$

$$\therefore \sum_x P_{X|A}(x) = \sum_x \frac{\mathbf{P}(\{X = x\} \cap A)}{\mathbf{P}(A)} = \frac{\sum_x \mathbf{P}(\{X = x\} \cap A)}{\mathbf{P}(A)} = \frac{\mathbf{P}(A)}{\mathbf{P}(A)} = 1$$

# Conditioning a Random Variable on an Event (2/2)

- A graphical illustration



**Figure 2.12:** Visualization and calculation of the conditional PMF  $p_{X|A}(x)$ . For each  $x$ , we add the probabilities of the outcomes in the intersection  $\{X = x\} \cap A$  and normalize by dividing with  $\mathbf{P}(A)$ .

$P_{X|A}(x)$  Is obtained by adding the probabilities of the outcomes that give rise to  $X = x$  and be long to the conditioning event  $A$

## Illustrative Examples (1/2)

- **Example 2.12.** Let  $X$  be the roll of a fair six-sided die and  $A$  be the event that the roll is an even number

$$\begin{aligned} P_{X|A}(x) &= \mathbf{P}(X = x | \text{roll is even}) \\ &= \frac{\mathbf{P}(X = x \text{ and } X \text{ is even})}{\mathbf{P}(X \text{ is even})} \\ &= \begin{cases} 1/3, & \text{if } x = 2, 4, 6 \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$

## Illustrative Examples (2/2)



- **Example 2.14.** A student will take a certain test repeatedly, up to a maximum of  $n$  times, each time with a probability  $p$  of passing, independently of the number of previous attempts.
  - What is the PMF of the number of attempts given that the student passes the test ?

Let  $X$  be a geometric random variable with parameter  $p$ , representing the number of attempts until the first success comes up

$$p_X(x) = (1-p)^{x-1} p$$

Let  $A$  be the event that the student pass the test within  $n$  attempts ( $A = \{X \leq n\}$ )

$$\therefore p_{X|A}(x) = \begin{cases} \frac{(1-p)^{x-1} p}{\sum_{m=1}^n (1-p)^{m-1} p}, & \text{if } x = 1, 2, \dots, n \\ 0, & \text{otherwise} \end{cases}$$





# Conditioning a Random Variable on Another (1/2)

- Let  $X$  and  $Y$  be two random variables associated with the same experiment. The conditional PMF  $p_{X|Y}$  of  $X$  given  $Y$  is defined as

$$p_{X|Y}(x|y) = \mathbf{P}(X = x|Y = y) = \frac{\mathbf{P}(X = x, Y = y)}{\mathbf{P}(Y = y)}$$

$$= \frac{p_{X,Y}(x, y)}{p_Y(y)}$$

$Y$  is fixed on some value  $y$

- Normalization Property  $\sum_x p_{X|Y}(x|y) = 1$

- The conditional PMF is often convenient for the calculation of the joint PMF

multiplication (chain) rule


$$p_{X,Y}(x, y) = p_Y(y)p_{X|Y}(x|y) (= p_X(x)p_{Y|X}(y|x))$$

# Conditioning a Random Variable on Another (2/2)

- The conditional PMF can also be used to calculate the marginal PMFs

$$p_X(x) = \sum_y p_{X,Y}(x,y) = \sum_y p_Y(y)p_{X|Y}(x|y)$$

- Visualization of the conditional PMF  $p_{X|Y}$



## An Illustrative Example (1/2)

- Example 2.14.** Professor May B. Right often has her facts wrong, and answers each of her students' questions incorrectly with probability  $1/4$ , independently of other questions. In each lecture May is asked 0, 1, or 2 questions with equal probability  $1/3$ .
  - What is the probability that she gives at least one wrong answer ?

Let  $X$  be the number of questions asked,

$Y$  be the number of questions answered wrong

$$\mathbf{P}(Y \geq 1) = \mathbf{P}(Y = 1) + \mathbf{P}(Y = 2)$$

$$= \mathbf{P}(X = 1, Y = 1) + \mathbf{P}(X = 2, Y = 1) + \mathbf{P}(X = 2, Y = 2)$$

$$\therefore \mathbf{P}(Y \geq 1) = \mathbf{P}(X = 1)\mathbf{P}(Y = 1|X = 1) + \mathbf{P}(X = 2)\mathbf{P}(Y = 1|X = 2)$$

$$+ \mathbf{P}(X = 2)\mathbf{P}(Y = 2|X = 2)$$

$$= \frac{1}{3} \cdot \frac{1}{4} + \frac{1}{3} \cdot \left[ \binom{2}{1} \frac{1}{4} \cdot \frac{3}{4} \right] + \frac{1}{3} \cdot \left[ \binom{2}{2} \frac{1}{4} \cdot \frac{1}{4} \right] = \frac{11}{48}$$


$$\binom{n}{k} p^k \cdot (1-p)^{n-k}$$

modeled as binomial distributions



## An Illustrative Example (2/2)

- Calculation of the joint PMF  $p_{X,Y}(x, y)$  in Example 2.14.



$y$			
2	0	0	1/48
1	0	4/48	6/48
0	16/48	12/48	9/48
	0	1	2
	$x$		

Joint PMF  $P_{X,Y}(x,y)$   
in tabular form

# Conditional Expectation

- Recall that a conditional PMF can be thought of as an ordinary PMF over a new universe determined by the conditioning event
- In the same spirit, a conditional expectation is the same as an ordinary expectation, except that it refers to the new universe, and all probabilities and PMFs are replaced by their conditional counterparts

# Summary of Facts About Conditional Expectations

- Let  $X$  and  $Y$  be two random variables associated with the same experiment
  - The conditional expectation of  $X$  given an event  $A$  with  $\mathbf{P}(A) > 0$ , is defined by

$$\mathbf{E} [X | A] = \sum_x x p_{X|A}(x)$$

- For a function  $g(X)$ , it is given by

$$\mathbf{E} [g(X) | A] = \sum_x g(x) p_{X|A}(x)$$

## Total Expectation Theorem (1/2)

- The conditional expectation of  $X$  given a value  $y$  of  $Y$  is defined by

$$\mathbf{E} [X | Y = y] = \sum_x xp_{X|Y}(x|y)$$

– We have

$$\mathbf{E} [X] = \sum_y p_Y(y) \sum_x xp_{X|Y}(x|y)$$

- Let  $A_1, \dots, A_n$  be disjoint events that form a partition of the sample space, and assume that  $P(A_i) > 0$ , for all  $i$ .

Then,

$$\mathbf{E} [X] = \sum_{i=1}^n \mathbf{P}(A_i) \mathbf{E} [X | A_i]$$

## Total Expectation Theorem (2/2)

- Let  $A_1, \dots, A_n$  be disjoint events that form a partition of an event  $B$ , and assume that  $P(A_i \cap B) > 0$ , for all  $i$ . Then,

$$\mathbf{E}[X | B] = \sum_{i=1}^n \mathbf{P}(A_i | B) \mathbf{E}[X | A_i \cap B]$$

- Verification of total expectation theorem

$$\begin{aligned} \mathbf{E}[X] &= \sum_x x p_X(x) = \sum_x x \sum_y p_{X,Y}(x,y) \\ &= \sum_x x \sum_y p_Y(y) p_{X|Y}(x|y) \\ &= \sum_y p_Y(y) \sum_x x p_{X|Y}(x|y) \\ &= \sum_y p_Y(y) \mathbf{E}[X | Y = y] \end{aligned}$$



## An Illustrative Example (1/2)

- Example 2.17.** Mean and Variance of the Geometric Random Variable

– A geometric random variable  $X$  has PMF  $p_X(x) = (1-p)^{x-1} p$ ,  $x=1,2,\dots$

Let  $A_1$  be the event that  $\{X = 1\}$

$A_2$  be the event that  $\{X > 1\}$

$$\mathbf{E}[X] = \mathbf{P}(A_1)\mathbf{E}[X|A_1] + \mathbf{P}(A_2)\mathbf{E}[X|A_2]$$

where


$$\mathbf{P}(A_1) = p, \mathbf{P}(A_2) = 1 - p \quad (??)$$

$$p_{X|A_1}(x) = \begin{cases} \frac{p}{p} = 1, & x = 1 \\ 0, & \text{otherwise} \end{cases}$$

$$p_{X|A_2}(x) = \begin{cases} (1-p)^{x-2} p \quad (??), & x > 1 \\ 0, & \text{otherwise} \end{cases}$$

Note that (See Example 2.13) :

$$p_{X|A}(x) = \begin{cases} \frac{(1-p)^{x-1} p}{\sum_{m=1}^n (1-p)^{m-1} p}, & \text{if } x = 1, 2, \dots, n \\ 0, & \text{otherwise} \end{cases}$$



$$\mathbf{E}[X|A_1] = 1 \cdot 1 + \sum_{x=2}^{\infty} x \cdot 0 = 1$$

$$\mathbf{E}[X|A_2] = 1 \cdot 0 + \sum_{x=2}^{\infty} x \cdot [(1-p)^{x-2} p]$$

$$= \sum_{x=2}^{\infty} x \cdot [(1-p)^{x-2} p]$$

$$= \sum_{x'=1}^{\infty} (x'+1)(1-p)^{x'-1} p$$

$$= \left[ \sum_{x'=1}^{\infty} x'(1-p)^{x'-1} p \right] + \left[ \sum_{x'=1}^{\infty} (1-p)^{x'-1} p \right]$$

$$= \mathbf{E}[X] + 1$$

$$\Rightarrow \mathbf{E}[X] = \mathbf{P}(A_1)\mathbf{E}[X|A_1] + \mathbf{P}(A_2)\mathbf{E}[X|A_2]$$

$$= \mathbf{P}(A_1) \cdot 1 + (1-p)(\mathbf{E}[X] + 1)$$

$$\therefore \mathbf{E}[X] = \frac{1}{p}$$

1

## An Illustrative Example (2/2)

$$\mathbf{E}[X^2] = \mathbf{P}(A_1)\mathbf{E}[X^2|A_1] + \mathbf{P}(A_2)\mathbf{E}[X^2|A_2]$$

$$\mathbf{E}[X^2|A_1] = 1^2 \cdot 1 + \sum_{x=2}^{\infty} x^2 \cdot 0 = 1$$

$$\begin{aligned} \mathbf{E}[X^2|A_2] &= 1^2 \cdot 0 + \sum_{x=2}^{\infty} x^2 \cdot (1-p)^{x-2} p \quad \curvearrowright \quad (\because x^2 = (x-1)^2 + 2x - 1) \\ &= \left[ \sum_{x=2}^{\infty} (x-1)^2 \cdot (1-p)^{x-2} p \right] + 2 \left[ \sum_{x=2}^{\infty} x \cdot (1-p)^{x-2} p \right] - \left[ \sum_{x=2}^{\infty} (1-p)^{x-2} p \right] \\ &= \left[ \sum_{x'=1}^{\infty} x'^2 \cdot (1-p)^{x'-1} p \right] + 2 \left[ \sum_{x=2}^{\infty} (x-1) \cdot (1-p)^{x-2} p \right] + 2 \left[ \sum_{x=2}^{\infty} (1-p)^{x-2} p \right] - \left[ \sum_{x=2}^{\infty} (1-p)^{x-2} p \right] \\ &= \mathbf{E}[X^2] + 2 \left[ \sum_{x'=1}^{\infty} x' \cdot (1-p)^{x'-1} p \right] + \left[ \sum_{x'=1}^{\infty} (1-p)^{x'-1} p \right] \quad (\text{set } x' = x - 1) \\ &= \mathbf{E}[X^2] + 2\mathbf{E}[X] + 1 \end{aligned}$$

$$\Rightarrow \mathbf{E}[X^2] = p \cdot 1 + (1-p)(\mathbf{E}[X^2] + 2\mathbf{E}[X] + 1)$$

$$\mathbf{E}[X^2] = \frac{1 + 2(1-p)\mathbf{E}[X]}{p} \quad \left( \text{we have shown that } \mathbf{E}[X] = \frac{1}{p} \right)$$

$$\mathbf{E}[X^2] = \frac{2}{p^2} - \frac{1}{p}$$

$$\therefore \text{var}(X) = \mathbf{E}[X^2] - (\mathbf{E}[X])^2 = \frac{1}{p^2} - \frac{1}{p} = \frac{1-p}{p^2}$$

# Independence of a Random Variable from an Event

- A random variable  $X$  is **independent of an event**  $A$  if

$$\mathbf{P}(X = x \text{ and } A) = \mathbf{P}(X = x)\mathbf{P}(A), \text{ for all } x$$

- Require two events  $\{X = x\}$  and  $A$  be independent for all  $x$
- If a random variable  $X$  is **independent of an event**  $A$  and  $\mathbf{P}(A) > 0$

$$\begin{aligned} p_{X|A}(x) &= \frac{\mathbf{P}(X = x \text{ and } A)}{\mathbf{P}(A)} \\ &= \frac{\mathbf{P}(X = x)\mathbf{P}(A)}{\mathbf{P}(A)} \\ &= \mathbf{P}(X = x) \\ &= p_X(x), \text{ for all } x \end{aligned}$$

## An Illustrative Example

- **Example 2.19.** Consider two independent tosses of a fair coin.
  - Let random variable  $X$  be the number of heads
  - Let random variable  $Y$  be 0 if the first toss is head, and 1 if the first toss is tail
  - Let  $A$  be the event that the number of head is even
    - Possible outcomes (T,T), (T,H), (H,T), (H,H)

$$p_X(x) = \begin{cases} 1/4, & \text{if } x = 0 \\ 1/2, & \text{if } x = 1 \\ 1/4, & \text{if } x = 2 \end{cases}$$

$$p_{X|A}(x) = \begin{cases} 1/2, & \text{if } x = 0 \\ 0, & \text{if } x = 1 \\ 1/2, & \text{if } x = 2 \end{cases}$$

$p_{X|A}(x) \neq p_X(x) \Rightarrow X$  and  $A$  are not independent!

$$p_Y(y) = \begin{cases} 1/2, & \text{if } y = 0 \\ 1/2, & \text{if } y = 1 \end{cases}$$

$$p_{Y|A}(y) = \frac{\mathbf{P}(Y = y \text{ and } A)}{\mathbf{P}(A)} = \begin{cases} 1/2, & \text{if } y = 0 \\ 1/2, & \text{if } y = 1 \end{cases}$$

$$\mathbf{P}(A) = 1/2$$

$p_{Y|A}(y) = p_Y(y) \Rightarrow Y$  and  $A$  are independent!

## Independence of a Random Variables (1/2)

- Two **random variables**  $X$  and  $Y$  are **independent** if

$$p_{X,Y}(x,y) = p_X(x)p_Y(y), \text{ for all } x, y$$

$$\text{or } \mathbf{P}(X = x, Y = y) = \mathbf{P}(X = x)\mathbf{P}(Y = y), \text{ for all } x, y$$

- If a random variable  $X$  is **independent of an random variable**  $Y$

$$p_{X|Y}(x|y) = p_X(x), \text{ for all } y \text{ with } p_Y(y) > 0 \text{ all } x$$

$$\begin{aligned} p_{X|Y}(x|y) &= \frac{p_{X,Y}(x,y)}{p_Y(y)} \\ &= \frac{p_X(x)p_Y(y)}{p_Y(y)} \\ &= p_X(x), \text{ for all } y \text{ with } p_Y(y) > 0 \text{ and all } x \end{aligned}$$

## Independence of a Random Variables (2/2)

- Random variables  $X$  and  $Y$  are said to be **conditionally independent**, given a positive probability event  $A$ , if

$$p_{X,Y|A}(x,y) = p_{X|A}(x)p_{Y|A}(y), \text{ for all } x, y$$

- Or equivalently,

$$p_{X|Y,A}(x|y) = p_{X|A}(x), \text{ for all } y \text{ with } p_{Y|A}(y) > 0 \text{ and all } x$$

- Note here that, as in the case of events, conditional independence may not imply unconditional independence and vice versa

## An Illustrative Example (1/2)

- **Figure 2.15:** Example illustrating that conditional independence may not imply unconditional independence
  - For the PMF shown, the random variables  $X$  and  $Y$  are not independent

- To show  $X$  and  $Y$  are not independent, we only have to find a pair of values  $(x, y)$  of  $X$  and  $Y$  that

$$p_{X|Y}(x|y) \neq p_X(x)$$

- For example,  $X$  and  $Y$  are not independent

$$p_{X|Y}(1|1) = 0 \neq p_X(1) = \frac{3}{20}$$

$y$	1	2	3	4
4	1/20	2/20	2/20	0
3	2/20	4/20	1/20	2/20
2	0	1/20	3/20	1/20
1	0	1/20	0	0

## An Illustrative Example (2/2)

- To show  $X$  and  $Y$  are not dependent, we only have to find all pair of values  $(x, y)$  of  $X$  and  $Y$  that

$$p_{X|Y}(x|y) = p_X(x)$$

- For example,  $X$  and  $Y$  are independent, conditioned on the event  $A = \{X \leq 2, Y \geq 3\}$


$$\mathbf{P}(A) = \frac{9}{20}, \quad p_{X|Y,A}(x|y) = \frac{\mathbf{P}(X = x \cap Y = y \cap A)}{\mathbf{P}(Y = y \cap A)}$$

$$p_{X|Y,A}(1|3) = \frac{2/20}{6/20} = \frac{1}{3}, \quad p_{X|A}(1) = \frac{3/20}{9/20} = 1/3$$

$$p_{X|Y,A}(1|4) = \frac{1/20}{3/20} = \frac{1}{3}$$

$$p_{X|Y,A}(2|3) = \frac{4/20}{6/20} = \frac{2}{3}, \quad p_{X|A}(2) = \frac{6/20}{9/20} = 2/3$$

$$p_{X|Y,A}(2|4) = \frac{2/20}{3/20} = \frac{2}{3}$$





# Functions of Two Independent Random Variables

- Given  $X$  and  $Y$  be two independent random variables, let  $g(X)$  and  $h(Y)$  be two functions of  $X$  and  $Y$ , respectively. Show that  $g(X)$  and  $h(Y)$  are independent.

Let  $U = g(X)$  and  $V = h(Y)$ , then

$$\begin{aligned} p_{U,V}(u,v) &= \sum_{\{(x,y)|g(x)=u, h(y)=v\}} p_{X,Y}(x,y) \\ &= \sum_{\{(x,y)|g(x)=u, h(y)=v\}} p_X(x) p_Y(y) \\ &= \sum_{\{x|g(x)=u\}} p_X(x) \sum_{\{y|h(y)=v\}} p_Y(y) \\ &= p_U(u) p_V(v) \end{aligned}$$

## More Factors about Independent Random Variables (1/2)

- If  $X$  and  $Y$  are independent random variables, then

$$\mathbf{E}[XY] = \mathbf{E}[X] \mathbf{E}[Y]$$

- As shown by the following calculation

$$\begin{aligned} \mathbf{E}[XY] &= \sum_x \sum_y xy \underline{p_{X,Y}(x,y)} \\ &= \sum_x \sum_y xy \underline{p_X(x)p_Y(y)} \quad \text{by independence} \\ &= \sum_x xp_X(x) \left[ \sum_y yp_Y(y) \right] \\ &= \mathbf{E}[X] \mathbf{E}[Y] \end{aligned}$$

- Similarly, if  $X$  and  $Y$  are independent random variables, then

$$\mathbf{E}[g(X)h(Y)] = \mathbf{E}[g(X)] \mathbf{E}[h(Y)]$$

## More Factors about Independent Random Variables (2/2)

- If  $X$  and  $Y$  are independent random variables, then

$$\text{var}(X + Y) = \text{var}(X) + \text{var}(Y)$$

- As shown by the following calculation

$$\begin{aligned} \text{var}(X + Y) &= \mathbf{E}\left[\left((X + Y) - \mathbf{E}[X + Y]\right)^2\right] \\ &= \mathbf{E}\left[(X + Y)^2 - 2(X + Y)(\mathbf{E}[X] + \mathbf{E}[Y]) + (\mathbf{E}[X] + \mathbf{E}[Y])^2\right] \\ &= \left[\sum_{x,y} (x + y)^2 p_{X,Y}(x, y)\right] - 2(\mathbf{E}[X] + \mathbf{E}[Y])\mathbf{E}[X] - 2(\mathbf{E}[X] + \mathbf{E}[Y])\mathbf{E}[Y] + \\ &\quad + (\mathbf{E}[X])^2 + 2 \cdot \mathbf{E}[X]\mathbf{E}[Y] + (\mathbf{E}[Y])^2 \\ &= \left[\sum_{x,y} x^2 p_{X,Y}(x, y)\right] + \left[\sum_{x,y} y^2 p_{X,Y}(x, y)\right] + 2\left[\sum_{x,y} xy p_{X,Y}(x, y)\right] \\ &\quad - (\mathbf{E}[X])^2 - (\mathbf{E}[Y])^2 - 2\mathbf{E}[X]\mathbf{E}[Y] \\ &= \left(\mathbf{E}[X^2] - (\mathbf{E}[X])^2\right) + \left(\mathbf{E}[Y^2] - (\mathbf{E}[Y])^2\right) = \text{var}(X) + \text{var}(Y) \end{aligned}$$

# More than Two Random Variables

- Independence of several random variables
  - Three random variable  $X$  ,  $Y$  and  $Z$  are independent if

$$p_{X,Y,Z}(x, y, z) = p_X(x)p_Y(y)p_Z(z) \text{ for all } x, y, z$$

? Compared to the conditions to be satisfied for three independent events  $A_1, A_2$  and  $A_3$  (in P.39 of the textbook)

- Any three random variables of the form  $f(X)$ ,  $g(X)$  and  $h(X)$  are also independent
- Variance of the sum of independent random variables
  - If  $X_1, X_2, \dots, X_n$  are independent random variables, then

$$\text{var}(X_1 + X_2 + \dots + X_n) = \text{var}(X_1) + \text{var}(X_2) + \dots + \text{var}(X_n)$$

## Illustrative Examples (1/3)

- **Example 2.20. Variance of the Binomial.** We consider  $n$  independent coin tosses, with each toss having probability  $p$  of coming up a head. For each  $i$ , we let  $X_i$  be the Bernoulli random variable which is equal to 1 if the  $i$ -th toss comes up a head, and is 0 otherwise.
  - Then,  $X = X_1 + X_2 + \cdots + X_n$  is a binomial random variable.

$$\therefore \text{var}(X_i) = p(1-p), \text{ for all } i$$

$$\therefore \text{var}(X) = \sum_{i=1}^n \text{var}(X_i) = np(1-p) \quad (\text{Note that } X_i \text{'s are independent!})$$

## Illustrative Examples (2/3)

- Example 2.21. Mean and Variance of the Sample Mean.** We wish to estimate the approval rating of a president, to be called B. To this end, we ask  $n$  persons drawn at random from the voter population, and we let  $X_i$  be a random variable that encodes the response of the  $i$ -th person:


$$X_i = \begin{cases} 1, & \text{if the } i\text{-th person approves B's performance} \\ 0, & \text{if the } i\text{-th person disapproves B's performance} \end{cases}$$

- Assume that  $X_i$  independent, and are the same random variable (Bernoulli) with the common parameter ( $p$  for Bernoulli), which is unknown to us

- $X_i$  are independent, and identically distributed (i.i.d.)

- If the sample mean  $S_n$  (is a random variable) is defined as

$$S_n = \frac{X_1 + X_2 + \dots + X_n}{n}$$



## Illustrative Examples (3/3)

- The expectation of  $S_n$  will be the true mean of  $X_i$

$$\begin{aligned}\mathbf{E}[S_n] &= \mathbf{E}\left[\frac{X_1 + X_2 + \cdots + X_n}{n}\right] \\ &= \frac{1}{n} \sum_{i=1}^n \mathbf{E}[X_i] \\ &= \mathbf{E}[X_i] \quad (= p \text{ for the Bernoulli we assumed here})\end{aligned}$$

- The variance of  $S_n$  will approximate 0 if  $n$  is large enough

$$\begin{aligned}\lim_{n \rightarrow \infty} \text{var}(S_n) &= \text{var}\left(\frac{X_1 + X_2 + \cdots + X_n}{n}\right) \\ &= \lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n \text{var}(X_i)}{n^2} = \lim_{n \rightarrow \infty} \frac{np(1-p)}{n^2} = \lim_{n \rightarrow \infty} \frac{p(1-p)}{n} = 0\end{aligned}$$

- Which means that  $S_n$  will be a good estimate of  $\mathbf{E}[X_i]$  if  $n$  is large enough

# Recitation

- SECTION 2.5 Joint PMFs of Multiple Random Variables
  - Problems 27, 28, 30
- SECTION 2.6 Conditioning
  - Problems 33, 34, 35, 37
- SECTION 2.6 Independence
  - Problems 42, 43, 45, 46